



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GFM80

Recombinant Mouse MIP-2 / CXCL2

Description

Macrophage Inflammatory Protein 2 (MIP-2), also known as CXCL2, is a small cytokine that is secreted by monocytes and neutrophils at sites of inflammation. MIP-2 functions through the chemokine receptor CXCR2 to act as a chemotactic agent for leukocytes and hematopoietic cells.

Length	73 aa
Molecular Weight	7.9 kDa
Source	E. coli
Accession Number	P10889
Purity	≥95% determined by reducing and non-reducing SDS-PAGE


Specifications

Alternative Names	Macrophage Inflammatory Protein 2, MIP2, CXCL2, chemokine (C-X-C motif) ligand 2, C-X-C motif chemokine 2
Biological Activity	Activity to be determined.
Endotoxin Level	≤1.00 EU/μg as measured by kinetic LAL
Formulation	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, pH 7.5
AA Sequence	AVVASELRQC CLKTLPRVDF KNIQSLSVTP PGPHCAQTEV IATLKGQKV CLDPEAPLVQ KIIQKILNKG KAN

Preparation and Storage

Reconstitution	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
Stability and Storage	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

Data



Non-reducing (-) and reducing (+) conditions in a 4 - 20% Tris-Glycine gel stained with Coomassie Blue. 1 μg of protein was loaded in each lane.
Mouse MIP-2 has a predicted Mw of 7.9 kDa.