



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

GFM84 Recombinant Mouse IL-10

Description

Interleukin-10 (IL-10) is an anti-inflammatory cytokine produced by macrophages and type 2 T helper (Th2) cells. IL-10 inhibits the production of pro-inflammatory cytokines such as interferon- γ (IFN- γ), Tumor Necrosis Factor α (TNF- α), interleukin-2 (IL-2), interleukin-3 (IL-3), interleukin-4 (IL-4), and Granulocyte-Macrophage Colony-Stimulating Factor (GM-CSF), made by macrophages and regulatory T cells. IL-10 also suppresses antigen presentation on antigen presenting cells, and enhances the survival, proliferation, and antibody production of B cells. Human IL-10 can act on mouse cells, while mouse IL-10 is not active on human cells.

Length	161 aa
Molecular Weight	18.9 kDa
Source	E. coli
Accession Number	P18893
Purity	$\geq 95\%$ determined by reducing and non-reducing SDS-PAGE


Specifications

Alternative Names	Interleukin-10, interleukin 10, IL10, IL 10, B-TCGF, CSIF, TGIF, cytokine synthesis inhibitory factor
Biological Activity	Mouse IL-10 is fully biologically active when compared to standard. The activity is determined by the ability to induce MC/9 cells proliferation and it is typically less than 5 ng/ml. This corresponds to an expected specific activity of 2.0×10^5 units/mg.
Endotoxin Level	≤ 1.00 EU/ μ g as measured by kinetic LAL
Formulation	Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 10 mM sodium phosphate, pH 7.5
AA Sequence	MSRGQYSRED NNCTHFPVGQ SHMLLELRTA FSQVKTFQ T KDQLDNILLT DSLMQDFKGY LGCQALSEMI QFYLVVEVMPQ AEKHGPEIKE HLNSLGEK LK TLRMLR RCH RFLPCENKSK AVEQVKSDFN KLQDQGVYKA MNEFDIFINC IEAYMMIKMK S

Preparation and Storage

Reconstitution	Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions.
Stability and Storage	12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed.

Data



Non-reducing (-) and reducing (+) conditions in a 4 - 20% Tris-Glycine gel stained with Coomassie Blue. 1 μ g of protein was loaded in each lane. Mouse IL-10 has a predicted Mw of 18.9 kDa.