



**SZABO
SCANDIC**

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic



GFR27

Recombinant Rat FGF-9

Description

Fibroblast Growth Factor 9 (FGF-9) is a mitogen and survival factor for nerve and mesenchymal cells. FGF-9 functions as an autocrine and paracrine factor to support the growth and survival of motor neurons and prostate tissue. FGF-9 expression in the gonad is also necessary for sex determination.

| | |
|-------------------------|---|
| Length | 207 aa |
| Molecular Weight | 23.4 kDa |
| Source | E. coli |
| Accession Number | P54130 |
| Purity | ≥95% determined by reducing and non-reducing SDS-PAGE |


Specifications

| | |
|----------------------------|---|
| Alternative Names | Fibroblast Growth Factor 9, FGF9, FGF 9, Glia Activating Factor, GAF, GAF2, heparin-binding growth factor 9, HBGF-9, SYNS3 |
| Biological Activity | Rat FGF-9 is fully biologically active when compared to standard. The activity is determined by the proliferation of NIH 3T3 cells. |
| Endotoxin Level | ≤1.00 EU/μg as measured by kinetic LAL |
| Formulation | Lyophilized from a sterile (0.2 micron) filtered aqueous solution containing 0.1% Trifluoroacetic Acid (TFA) |
| AA Sequence | MPLGEVGSYFG VQDAVPFGNV PVLPVDSPLV LNDHLGQSEA GGLPRGPAVT DLDHLKGILR RRQLYCRTGF HLEIFPNTI QGTRKDHSRF GILEFISIAV GLVSIRGVDS GLYLGGMNEKG ELYGSEKLTQ ECVFREQFEE NWYNTYSSNL YKHVDTGRYY YVALNKDGTP REGTRTKRHQ KFTHFLPRPV DPDKVPELYK DILSQS |

Preparation and Storage

| | |
|------------------------------|---|
| Reconstitution | Centrifuge vial before opening. When reconstituting the product, gently pipet and wash down the sides of the vial to ensure full recovery of the protein into solution. It is recommended to reconstitute the lyophilized product with sterile water at 0.1 mg/ml, which can be further diluted into other aqueous solutions. |
| Stability and Storage | 12 months from date of receipt when stored at -20°C to -80°C as supplied. 1 month when stored at 4°C after reconstituting as directed. 3 months when stored at -20°C to -80°C after reconstituting as directed. |

Data



Induced proliferation of NIH 3T3 cells assay for Rat FGF-9. Cell proliferation was measured to calculate the ED50.