



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## PPH317 PODS<sup>®</sup> Human NGF (Mature)

### Description

The product contains the polyhedrin protein co-crystallized with Human NGF (Mature). Nerve Growth Factor (NGF) is a neurotrophic factor that is important for the development and maintenance of sensory and sympathetic neurons. NGF signals through the low affinity nerve growth factor receptor (LNGFR) and the tropomyosin receptor kinase A (TrkA) to activate PI3K, Ras, and PLC signaling pathways. NGF is also involved in the growth, differentiation, and survival of B lymphocytes.

|                         |   |
|-------------------------|---|
| <b>Length</b>           | 268 aa  |
| <b>Molecular Weight</b> | 60.2 kDa  |
| <b>Source</b>           | <i>Spodoptera frugiperda (Sf9) cell culture</i> |
| <b>Accession Number</b> | P01138  |

### Usage Recommendation

PODS<sup>®</sup> co-crystals provide a depot of proteins which are steadily secreted. It has been estimated that the biological activity of 50 million PODS<sup>®</sup> co-crystals generates the same peak dose as 3.3 µg of standard recombinant protein. However, at 5 days following the start of seeding the PODS<sup>®</sup> co-crystals, there are more than 50% of these peak levels still present in the culture system. Ultimately, the amount of PODS<sup>®</sup> co-crystals that is optimal for a particular experiment should be determined empirically. Based on previous data, we suggest using 50 million PODS<sup>®</sup> co-crystals in place of 3.3 µg of standard growth factor as a starting point. To control for cross-reactivity with cells or as a negative control, we recommend using PODS<sup>®</sup> growth factors alongside [PODS<sup>®</sup> Empty crystals](http://www.cellgs.com/products/podsand8482-empty.html), as the latter do not contain or release cargo protein.

### Specifications

|                          |   |
|--------------------------|---|
| <b>Alternative Names</b> | Migration Inhibitory Factor, GIF, phenylpyruvate tautomerase, glycosylation-inhibiting factor, L-dopachrome tautomeras  |
| <b>Endotoxin Level</b>   | <0.06 EU/ml as measured by gel clot LAL assay   |
| <b>Formulation</b>       | PODS <sup>®</sup> were lyophilized from a volatile solution   |
| <b>AA Sequence</b>       | MADVAGTSNR DFRGREQRLF NSEQYNNNS KNSRPSTSLY KKAGSSSSHP IFHRGEFSVC<br>DSVSVWVGDK TTATDIKKE VMVLGEVNIN NSVFKQYFFE TKCRDPNPVD SGCRGIDSKH<br>WNSYCTTHT FVKALTMGK QAAWRFIRID TACVCVLSRK AVRRA |

### Preparation and Storage

|                              |  |
|------------------------------|--|
| <b>Reconstitution</b>        | PODS <sup>®</sup> co-crystals may be reconstituted at 200 million co-crystals/ml in water. 20% glucose has a buoyant density closer to PODS <sup>®</sup> co-crystals and can be useful for aliquoting. PODS <sup>®</sup> co-crystals are highly stable when stored in aqueous solution (pH range 6 - 8). |
| <b>Stability and Storage</b> | Upon receipt, store at 4°C. PODS <sup>®</sup> co-crystals are stable for at least 1 year when dry and 6 months when resuspended.   |