



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

PPH77 PODS[®] Human IFN- γ

Description

The product contains the polyhedrin protein co-crystalized with Human IFN- γ . Interferon γ (IFN- γ) is a type II interferon that is critical during adaptive and innate immune responses to infection. IFN- γ is produced by T cells and Natural Killer cells following antigen-specific activation. IFN- γ binds IFN- γ receptors (IFN- γ R1 and IFN- γ R2), which are expressed on most immune cells, to activate the JAK-STAT pathway. IFN- γ induced signaling increases the expression of class 1 major histocompatibility complex (MHC) molecules. Human IFN- γ is not cross-reactive with mouse IFN- γ .

Length	211 aa
Molecular Weight	49.4 kDa
Source	<i>Spodoptera frugiperda (Sf9) cell culture</i>
Accession Number	P01579

Usage Recommendation

PODS[®] co-crystals provide a depot of proteins which are steadily secreted. It has been estimated that the biological activity of 50 million PODS[®] co-crystals generates the same peak dose as 3.3 μ g of standard recombinant protein. However, at 5 days following the start of seeding the PODS[®] co-crystals, there are more than 50% of these peak levels still present in the culture system. Ultimately, the amount of PODS[®] co-crystals that is optimal for a particular experiment should be determined empirically. Based on previous data, we suggest using 50 million PODS[®] co-crystals in place of 3.3 μ g of standard growth factor as a starting point. To control for cross-reactivity with cells or as a negative control, we recommend using PODS[®] growth factors alongside [PODS[®] Empty crystals](http://www.cellgs.com/products/podsand8482-empty.html), as the latter do not contain or release cargo protein.

Specifications

Alternative Names	Interferon- γ , interferon-gamma, interferon gamma, type II interferon, T cell interferon, immune Interferon, MAF, IFG, IFI, IFN-gamma, IFN gamma
Endotoxin Level	<0.06 EU/ml as measured by gel clot LAL assay
Formulation	PODS [®] were lyophilized from a volatile solution
AA Sequence	MQDPYVKEAE NLKKYFNAGH SDVADNGTLF LGILKNWKEE SDRKIMQSQI VSFYFKLFFKN FKDDQSIQKS VETIKEDMNV KFFNSNKKKR DDFEKLTNYS VTDLNVQRKA IHELIQVMAE LSPAAGTKGR KRSQMLFRGR RASQDPAFLY KVVVDGYLLAF NSQRRSHTLR LLGPFQYFNF SETDRGHPLF RLPLKYPSKA IPADELIDNL H

Preparation and Storage

Reconstitution	PODS [®] co-crystals may be reconstituted at 200 million co-crystals/ml in water. 20% glucose has a buoyant density closer to PODS [®] co-crystals and can be useful for aliquoting. PODS [®] co-crystals are highly stable when stored in aqueous solution (pH range 6 - 8).
Stability and Storage	Upon receipt, store at 4°C. PODS [®] co-crystals are stable for at least 1 year when dry and 6 months when resuspended.