



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



Human versican/PG-M/PG-350 ELISA kit

Product Code	CSB-E11884h
Abbreviation	VCAN
Uniprot No.	P13611
Product Type	ELISA Kit
Immunogen Species	Homo sapiens (Human)
Sample Types	serum, plasma, tissue homogenates.
Detection Range	0.04 ng/ml-30 ng/ml.
Sensitivity	0.04 ng/ml.
Assay Time	1-5h
Sample Volume	50-100ul
Detection Wavelength	450 nm
Lead Time	3-5 working days
Research Area	Signal Transduction
Gene Names	VCAN
Tag Info	quantitative
Protein Description	Sandwich

Description

The human VCAN ELISA kit (CSB-E11884h) is designed for the quantitative measurement of human VCAN protein in samples. It shows excellent specificity for human VCAN. It uses the bi-antibody sandwich enzyme immunoassay technique. This assay employs a biotin-conjugated VCAN antibody that recognizes the analyte bound by the immobilized VCAN antibody, forming an antibody-analyte-antibody complex. The immune complex is further detected by avidin-conjugated HRP. The TMB solution is added into the wells and turns blue and finally turns yellow after the addition of the stop solution. Solution color develops in proportion to the amount of VCAN in the sample. The O.D. value is measured using a microplate reader at 450 nm and is used to determine the concentration of the human VCAN in the sample.

VCAN is a large extracellular matrix (ECM) proteoglycan that is significantly increased expressed and accumulated during inflammation in most diseases. It plays an essential role during development and is an important component of tissue inflammation caused by infection and tissue injury. VCAN interacts with its receptors such as CD44, PSGL-1, and TLR2, initiating a signaling cascade that affects the phenotypes of immune and inflammatory cells. It is also implicated in multiple processes of cancer development, including proliferative signaling, evasion of growth-suppressor pathways, modulation of cell death, promotion of neoangiogenesis, and tissue invasion & metastasis.



Product Precision

Linearity

Recovery