



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com


[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



VN1R1 Antibody

Product Code	CSB-PA861973LA01HU
Abbreviation	Vomer nasal type-1 receptor 1
Storage	Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.
Uniprot No.	Q9GZP7
Immunogen	Recombinant Human Vomer nasal type-1 receptor 1 protein (1-56AA)
Raised In	Rabbit
Species Reactivity	Human
Tested Applications	ELISA, WB, IF; Recommended dilution: WB:1:500-1:5000, IF:1:50-1:200
Relevance	Putative pheromone receptor.
Form	Liquid
Conjugate	Non-conjugated
Storage Buffer	Preservative: 0.03% Proclin 300 Constituents: 50% Glycerol, 0.01M PBS, pH 7.4
Purification Method	>95%, Protein G purified
Isotype	IgG
Clonality	Polyclonal
Alias	Vomer nasal type-1 receptor 1 (G-protein coupled receptor GPCR24) (hGPCR24) (V1r-like receptor 1) (V3r-related gene protein) (Vomer nasal olfactory receptor chromosome 19 subtype I member 1), VN1R1, V1RL1, VNR1911
Species	Homo sapiens (Human)
Research Area	Others
Target Names	VN1R1

Image



Western Blot

Positive WB detected in: HeLa whole cell lysate, LO2 whole cell lysate, A549 whole cell lysate, PC-3 whole cell lysate


All lanes: VN1R1 antibody at 3.3µg/ml

Secondary

Goat polyclonal to rabbit IgG at 1/50000 dilution

Predicted band size: 41 kDa

Observed band size: 41 kDa



Immunofluorescence staining of HeLa cells with CSB-PA861973LA01HU at 1:133, counter-stained with DAPI. The cells were fixed in 4% formaldehyde, permeabilized using 0.2% Triton X-100 and blocked in 10% normal Goat Serum. The cells were then incubated with the antibody overnight at 4°C. The secondary antibody was Alexa Fluor 488-conjugated AffiniPure Goat Anti-Rabbit IgG(H+L).