



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Prdm9 Polyclonal Antibody

Catalog Number: E-AB-92241



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.


## Description

<b>Reactivity</b>	Mouse, Rat
<b>Immunogen</b>	Recombinant fusion protein of mouse Prdm9
<b>Host</b>	Rabbit
<b>Isotype</b>	IgG
<b>Purification</b>	Affinity purification
<b>Conjugation</b>	Unconjugated
<b>Formulation</b>	PBS with 0.05% proclin300, 50% glycerol, pH7.3.

## Applications Recommended Dilution

<b>WB</b>	1:500-1:2000
-----------	--------------


## Data



Western blot analysis of extracts of various cell lines using Prdm9 Polyclonal Antibody at 1:1000 dilution.

**Observed Mw:100KDa**

**Calculated Mw:97kDa**



Western blot analysis of extracts of various cell lines using Prdm9 Polyclonal Antibody at 1:1000 dilution.

## Preparation & Storage

**Storage** Store at -20°C. Avoid freeze/thaw cycles.

## Background

Histone methyltransferase that sequentially mono-, di-, and tri-methylates both 'Lys-4' (H3K4 and 'Lys-36' (H3K36) of histone H3 to produce respectively trimethylated 'Lys-4' (H3K4me3) and trimethylated 'Lys-36' (H3K36me3) histone H3 and plays a key role in meiotic prophase by determining hotspot localization thereby promoting meiotic recombination. Also can methylate all four core histones with H3 being the best substrate and the most highly modified. Is also able, on one hand, to mono and di-methylate H4K20 and on other hand to trimethylate H3K9 with the di-methylated H3K9 as the best substrate. During meiotic prophase, binds specific DNA sequences through its zinc finger domains thereby determining hotspot localization where it promotes local H3K4me3 and H3K36me3 enrichment on the same nucleosomes through its histone methyltransferase activity. Thereby promotes double-stranded breaks (DSB) formation, at this subset of PRDM9-binding sites, that initiates meiotic recombination for the proper meiotic progression.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017