

### Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

#### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

#### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

## **Elabscience**®

### Elab Fluor<sup>®</sup> Red 780 Anti-Human CD127/IL-7RA Antibody[A019D5]

Catalog No.E-AB-F1152SStorageStore at 2~8°C, Avoid freeze / thaw cycles

ReactivityHumanApplicationsFCM

**Important Note:** Centrifuge before opening to ensure complete recovery of vial contents.

#### **Antigen Information**

Alternate Names	Interleukin-7 receptor subunit alpha,IL-7RA,CDw127,CD127,IL-7Rα
Alter nate Maines	interedkin-7 receptor subunit alpha,iL-7KA,CDw127,CD127,iL-7KG
Uniprot ID	P16871
Gene ID	3575
Background	CD127 is a 60-90 kD type I transmembrane glycoprotein also known as IL-7 receptor $\alpha$ chain or
	IL-7R $\alpha$ . It forms a heterodimer with the common $\gamma$ chain ( $\gamma c$ or CD132) which is shared with the
	receptors for IL-2, IL-4, IL-9, IL-13, IL-15, and IL-21. CD127 is expressed on immature B cells
	through early pre-B stage cells, thymocytes (except CD4/CD8 double positive thymocytes),
	peripheral T cells, and bone marrow stromal cells. CD127 has been reported to be a useful marker
	for identifying memory and effector T cells. Studies have shown that CD127 expression is down-
	modulated on Treg cells. It can be used as a marker for differentiation of Treg and conventional T
	cells. The ligation of IL-7 with its receptor is important for stimulation of mature and immature T
	cells as well as immature B cell proliferation and development.

#### **Product Details**

Form	Liquid
Size	20Tests/100Tests/100Tests×2
Clone No.	A019D5
Host	Mouse
Isotype	Mouse IgG1, ĸ
Reactivity	Human
Application	FCM
Isotype Control	Elab Fluor <sup>®</sup> Red 780 Mouse IgG1, κ Isotype Control[MOPC-21] [Product E-AB-F09792S]
Storage Buffer	Phosphate buffered solution, pH 7.2, containing 0.09% stabilizer and 1% protein protectant.
Shipping	Biological ice pack at 4 °C
Stability & Storage	Keep as concentrated solution.
	Store at 2~8°C and protected from prolonged exposure to light.Do not freeze.
	This product is guaranteed up to one year from purchase.

**For Research Use Only** 

# Elabscience®

#### Fluorophore


Conjugation: Elab Fluor<sup>®</sup> Red 780

Elab Fluor<sup>®</sup> Red 780 is designed to be excited by the Red (627-640 nm) laser and detected using an optical filter centered near 770 nm (e.g., a 780/60 nm bandpass filter).

#### **Recommended usage**

Each lot of this antibody is quality control tested by flow cytometric analysis. The amount of the reagent is suggested to be used 5  $\mu$ L of antibody per test (million cells in 100  $\mu$ L staining volume or per 100  $\mu$ L of whole blood). Please check your vial before the experiment. Since applications vary, the appropriate dilutions must be determined for individual use.

#### **Product data**



Human peripheral blood lymphocytes are stained with FITC Anti-Human CD3 Antibody and Elab Fluor<sup>®</sup> Red 780 Anti-Human CD127/IL-7RA Antibody (Left). Lymphocytes are stained with FITC Anti-Human CD3 Antibody and Elab Fluor<sup>®</sup> Red 780 Mouse IgG1, κ Isotype Control (Right).

#### **Related Information**

- 1. Sample Preparation for Flow Cytometry https://www.elabscience.com/List-detail-5594.html
- 2. Staining Cell Surface Targets for Flow Cytometry https://www.elabscience.com/List-detail-5568.html
- 3. Flow Cytometry Troubleshooting Tips <u>https://www.elabscience.com/List-detail-5593.html</u>
- 4. How to select the appropriate detection channel through the spectrogram? <u>https://www.elabscience.com/List-detail-459742.html</u>