



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

(FOR RESEARCH USE ONLY. DO NOT USE IT IN CLINICAL DIAGNOSIS !)

10× ACK Lysis Buffer

Catalog No: E-CK-A105

Size: 100 mL / 200 mL / 500 mL

Cat.	Products	100 mL	200 mL	500 mL	Storage
E-CK-A105	10× ACK Lysis Buffer	100 mL	200 mL	500 mL	2~8°C
Manual			One Copy		

This manual must be read attentively and completely before using this product.

If you have any problems, please contact our Technical Service Center for help.

Phone: 240-252-7368(USA) Fax: 240-252-7376(USA)

Email: techsupport@elabscience.com

Website: www.elabscience.com

Please kindly provide us the lot number (on the outside of the box) of the kit for more efficient service.

Introduction

Red Blood Cell Lysis Buffer (RBC Lysis), also known as ACK Lysis Buffer, is a kind of solution used to lyse and remove enucleated red blood cells from blood or tissue samples of human or mouse.

This product is sterile. After treating with this buffer, the single cell suspensions can be used for subsequent primary culture, cell fusion, extraction of nucleic acid or protein, and various routine analysis and detection.

Instructions

10×ACK Lysis Buffer is concentrated, dilute with ddH₂O to 1×working solution before use.

For example: take 1 mL 10×ACK Lysis Buffer [E-CK-A105], add to 9 mL ddH₂O. The mixture is 1×ACK Lysis Working Buffer.

Experimental Procedure

➤ For Mouse Spleen or Bone Marrow Samples

1. Centrifuge single cell suspension of the sample at 300 g for 5 min, discard the supernatant.
2. Add 2 mL of 1 ×ACK Lysis Working Buffer to the sample to resuspend the cells and incubate at RT for 2~3 min (Incubation time should be adjusted accordingly to the actual situation).
3. Add 10 mL PBS to stop the lysis reaction.
4. Centrifuge at 300 g for 5 min and discard the supernatant.
5. Resuspend the cells with PBS or similar physiological buffer for subsequent experiments.

➤ For Human Peripheral Blood Samples

1. Add 100 µL fresh blood into a centrifuge tube, add 2 mL of 1 ×ACK Lysis Working Buffer and mix thoroughly.
2. Put the centrifuge tube on ice for 5 min, until the fresh blood becomes transparent.
3. Centrifuge at 300 g for 5 min and discard the supernatant. Use a pipette to remove the residual supernatant carefully.
4. Add 2 mL PBS to the tube and repeat step 3.
5. Resuspend the cells with PBS or similar physiological buffer for subsequent experiments.

Storage

Store at 2~8 °C. Avoid of freezing. Valid for 12 months.

Cautions

1. This product is sterile. Please open and use it in laminar flow bench.
2. This reagent is for research use only.
3. For your safety and health, please wear the lab coat and disposable gloves before the experiments.