



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

0.25% Trypsin Solution

Cat. No:	E-EL-SR001
Size:	100mL

General Information

Product Form	Liquid
Concentration	100mM
Phenol red	negative
Dissolvent	10mM PBS
EDTA	0.2g/L
Trypsin	2.5g/L
Storage	-20 °C, shading light
Shipping	Ice bag
Expiration date	12 months

Background

Trypsin is a serine hydrolase that can cut the base-side segment of the lysine and arginine residue in the polypeptide chain, hydrolyze the protein between the cells, and destroy the connection between the cells, so that the tissue or the adherent cells are dispersed into a single cell. The activity of the trypsin is related to the characteristics of the tissue or the cells, the concentration of the trypsin, the temperature and the time of action. At pH 8.0 and 37 °C, the effect of the trypsin is the best. Therefore, the concentration of trypsin, the temperature and the time of the pancreatic enzyme should be grasped to avoid the excessive damage of the cells. In general, the frequently-used working concentration of the trypsin is 0.25%, while the concentration of trypsin for the semi-adherent cells or the cells is 0.05%. Since EDTA is able to bind Ca²⁺ and Mg²⁺ to destroy the cell connection to promote the dissociation of the cells, a certain amount of EDTA is often added to the trypsin solution to enhance the dissociation effect.

Notes

1. This product is for research use only.
2. It is necessary to pay attention to the aseptic operation and avoid the pollution.
3. It is not suitable for long time storage at room temperature or 4 °C.
4. Thaw in 2 -8 °C and mix fully before use. Avoid repeated freezing and thawing. It is recommended to aliquot the solution and store at -20 °C.
5. Because the sensitivity to trypsin of different tissues or cells are differently. The reaction time of trypsin should be determined according to the actual situation. Otherwise it will affect the the adhesion and growth of cells.