

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





Penicillin-Streptomycin-Glutamine Solution (100×)

Cat. No: PB180122 Size: 100mL

General Information

Product Form Liquid Concentration $100 \times$ pH 6.1-6.4 L-glutamine 200mM

Dissolvent 10mM PBS (pH6.2)

Streptomycin sulfate 10mg/mL Penicillin G 10kU/mL

Antimicrobial spectrum Gram-positive bacteria & Gram-negative

bacteria

Storage -20°C, Shading light Shipping Ice bag or dry ice

Expiration date 12 months

Background

Penicillin-streptomycin solution mixture is the most commonly used antibiotic to prevent microbial contamination in vitro. Penicillin can interfere with the synthesis of bacterial cell wall, especially for Gram-positive bacteria. Streptomycin could bind to ribosomal ribosome for 30S and inhibit the synthesis of bacterial protein. It was effective for Gram-negative bacteria and Gram-positive bacteria, but especially for Gram-negative bacteria. The combined use of penicillin and streptomycin can prevent most bacterial contamination, but penicillin solution is sensitive to temperature and pH, easy to degrade at room temperature, and needs cryopreservation, the stability of penicillin solution is the most stable when pH is 6.0 - 6.5. Streptomycin was relatively stable, and pH 5.0-7.5 was the most stable. L-glutamine is an essential element in cell culture that is not only an energy source of the cells, but also an important substance involved in protein synthesis and nucleic acid metabolism. The deficiency of L-glutamine will lead to poor cell growth and even death. L-glutamine is unstable in solution and easy to degrade spontaneously, so it is necessary to supplement a certain amount of L-glutamine in culture medium to meet the needs of cell growth after a period of time or when cells grow badly.

Notes

- 1. This product is for research use only.
- 2. This product is sterilized by 0.22μ m filtration.
- 3. It is necessary to pay attention to the aseptic operation and avoid the pollution.
- 4. Avoid repeated freezing and thawing. It is recommended to defrost at 2-8°C and aliquot the solution and store at -20°C.
- 5. This product is a concentrated solution. Please dilute it as needed.