



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## 50mg/mL G-418 Solution

Cat. No: PB180125  
Size: 10mL

### General Information

Product Form	Liquid
Concentration	50mg/mL
pH	7.2-7.4
Recommended working concentration	50mg/mL
Dissolvent	10mM HEPES
Storage	-20°C, Shading light
Shipping	Ice bag or dry ice
Expiration date	24 months

### Background

G418, an aminoglycoside antibiotic, is the most commonly used resistance screening reagent for stable transfection in molecular genetic experiments. G418 can interfere with the function of 80S ribosomes and affect protein synthesis in eukaryotic cells. It is mainly used in mammalian, plant or yeast cell selection. G418 is toxic to bacteria, yeast, higher plants, mammalian cells, protozoa, and worms. Its resistance gene is a dominant phenotype, and it is located on Tn601 (903) and Tn5 transposons. Although this gene is derived from bacteria, it can also be expressed in eukaryotic cells. The introduction of resistance genes into cells results in resistance to G418, allowing cells to grow in a culture medium containing G418. This selective feature of G418 has been widely used in gene transfer, gene knockout, resistance screening and transgenic animals.

### Notes

1. This product is for research use only.
2. This product is sterilized by 0.22 μ m filtration.
3. It is necessary to pay attention to the aseptic operation and avoid the pollution.
4. Avoid repeated freezing and thawing. It is recommended to defrost at 2-8°C and aliquot the solution and store at -20°C.