



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

Locke's Solution (low calcium)

Cat. No: PB180344
Size: 500mL

General Information

Product Form	Liquid
Concentration	1×
pH	7.2-7.4
D-glucose	2000mg/L
NaHCO ₃	200mg/L
Phenol red	Negative
CaCl ₂ (anhydrous)	50mg/L
Storage	2-30°C
Shipping	RT
Expiration date	18 months

Background

Balanced Salt Solution (Physiological Solution) have the properties of buffer solution (regulate pH), normal saline (maintain osmotic pressure) and culture medium (provide nutrition). It can meet the basic needs of survival and metabolism of tissues, organs or cells in vitro. A small amount of phenolic red was added to some equilibrium salt solutions to indicate the pH change of the solution. Locke's solution is a kind of balanced salt solution and the main components are NaCl, KCl, NaH₂PO₄, CaCl₂, NaHCO₃ and glucose. Locke's solution is commonly used in mammalian heart, uterus and other isolated organs. When used for irrigation, oxygen must be infused for 15 minutes before use. Locke's solution with low content calcium is used for perfusion of isolated small intestine and isolated organs of guinea pig.

Notes

1. This product is for research use only.
2. This product is sterilized by 0.22 μ m filtration.
3. It is necessary to pay attention to the aseptic operation and avoid the pollution.
4. Since the solution contains calcium, there maybe precipitation produced, which will be dissolved after 37°C water bath incubation.