

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Recombinant Human Parvalbumin Alpha/PVALB Protein (His Tag)



Catalog Number:PDEH100006

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Parvalbumin Alpha;PVALB

SpeciesHumanExpression HostE.coliSequenceSer2-Ser110AccessionP20472

Calculated Molecular Weight13.1 kDaObserved molecular weight14 kDaTagC-His

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin < 1.0 EU per µg of the protein as determined by the LAL method.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.Formulation Lyophilized from a 0.2 μm filtered solution of 20mM PB, 150mM NaCl, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed man

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Parvalbumin α (PVALB) is a member of the parvalbumin family. PVALB is a high affinity calcium ion-binding protein, with two EF hand domains. PVALB is structurally and functionally similar to calmodulin and troponin C, it can bind two calcium ions. Parvalbumin is thought to be involved in relaxation after contraction in muscle. Parvalbumin is expressed in a specific population of GABAergic interneurons, which are believed to have a role in maintaining the balance between excitation and inhibition in the cortex as well as the hippocampus.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com