

# Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

# Recombinant Human IL-6 Protein (His Tag)

Catalog Number:PDEH100050



Note: Centrifuge before opening to ensure complete recovery of vial contents.

ED50 for this effect is 0.27 to 0.81 ng/mL.         Properties         Purity       > 90 % as determined by reducing SDS-PAGE.         Endotoxin       Please contact us for more information.	Description	
Expression HostE.coliSequenceVal30-Met212AccessionP05231Calculated Molecular Weight20.8 kDaObserved molecular weight25-28 kDaTagN-HisBioactivityMeasured in a cell proliferation assay using TF-1 human erythroleukemic cells. The ED50 for this effect is 0.27 to 0.81 ng/mL.PropertiesPurity> 90 % as determined by reducing SDS-PAGE.EndotoxinPlease contact us for more information.StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from sterile 50mM Tris-HCl, pH 8.0 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.	Synonyms	Interleukin-6;IL-6;B-cell hybridoma growth factor;Interleukin HP-1
SequenceVal30-Met212AccessionP05231Calculated Molecular Weight20.8 kDaObserved molecular weight25-28 kDaTagN-HisBioactivityMeasured in a cell proliferation assay using TF-1 human erythroleukemic cells. The ED50 for this effect is 0.27 to 0.81 ng/mL.PropertiesPurity> 90 % as determined by reducing SDS-PAGE.EndotoxinPlease contact us for more information.StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from sterile 50mM Tris-HCl, pH 8.0 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.ReconstitutionPlease refer to the printed manual for detailed information.	Species	Human
AccessionP05231Calculated Molecular Weight20.8 kDaObserved molecular weight25-28 kDaTagN-HisBioactivityMeasured in a cell proliferation assay using TF-1 human erythroleukemic cells. The ED50 for this effect is 0.27 to 0.81 ng/mL.PropertiesPurity> 90 % as determined by reducing SDS-PAGE.EndotoxinPlease contact us for more information.StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Expression Host	E.coli
Calculated Molecular Weight20.8 kDaObserved molecular weight25-28 kDaTagN-HisBioactivityMeasured in a cell proliferation assay using TF-1 human erythroleukemic cells. The ED50 for this effect is 0.27 to 0.81 ng/mL.PropertiesPurity> 90 % as determined by reducing SDS-PAGE.EndotoxinPlease contact us for more information.StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Sequence	Val30-Met212
Observed molecular weight25-28 kDaTagN-HisBioactivityMeasured in a cell proliferation assay using TF-1 human erythroleukemic cells. The ED50 for this effect is 0.27 to 0.81 ng/mL.PropertiesPurity> 90 % as determined by reducing SDS-PAGE.EndotoxinPlease contact us for more information.StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at <-20°C for 3 months.	Accession	P05231
TagN-HisBioactivityMeasured in a cell proliferation assay using TF-1 human erythroleukemic cells. The ED50 for this effect is 0.27 to 0.81 ng/mL.PropertiesPurity> 90 % as determined by reducing SDS-PAGE.EndotoxinPlease contact us for more information.StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Calculated Molecular Weight	20.8 kDa
Bioactivity       Measured in a cell proliferation assay using TF-1 human erythroleukemic cells. The ED50 for this effect is 0.27 to 0.81 ng/mL.         Properties         Purity       > 90 % as determined by reducing SDS-PAGE.         Endotoxin       Please contact us for more information.         Storage       Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.         Shipping       This product is provided as lyophilized powder which is shipped with ice packs.         Formulation       Lyophilized from sterile 50mM Tris-HCl, pH 8.0 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.         Reconstitution       Please refer to the printed manual for detailed information.	Observed molecular weight	25-28 kDa
ED50 for this effect is 0.27 to 0.81 ng/mL.         Properties         Purity       > 90 % as determined by reducing SDS-PAGE.         Endotoxin       Please contact us for more information.         Storage       Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.         Shipping       This product is provided as lyophilized powder which is shipped with ice packs.         Formulation       Lyophilized from sterile 50mM Tris-HCl, pH 8.0 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.         Reconstitution       Please refer to the printed manual for detailed information.	Tag	N-His
Purity> 90 % as determined by reducing SDS-PAGE.EndotoxinPlease contact us for more information.StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.ShippingThis product is provided as lyophilized powder which is shipped with ice packs.FormulationLyophilized from sterile 50mM Tris-HCl, pH 8.0 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.ReconstitutionPlease refer to the specific buffer information in the printed manual.	Bioactivity	Measured in a cell proliferation assay using TF-1 human erythroleukemic cells. The ED50 for this effect is 0.27 to 0.81 ng/mL.
EndotoxinPlease contact us for more information.StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Properties	
StorageGenerally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Purity	> 90 % as determined by reducing SDS-PAGE.
-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	Endotoxin	Please contact us for more information.
Formulation       Lyophilized from sterile 50mM Tris-HCl, pH 8.0 Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.         Reconstitution       Please refer to the printed manual for detailed information.	Storage	
Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.ReconstitutionPlease refer to the printed manual for detailed information.	Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
•	Formulation	Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.
Data	Reconstitution	Please refer to the printed manual for detailed information.
	Data	



> 90 % as determined by reducing SDS-PAGE.

#### Background

Interleukin-6 (IL-6) is a pleiotropic, alpha -helical, 22-28 kDa phosphorylated and variably glycosylated cytokine that plays important roles in the acute phase reaction, inflammation, hematopoiesis, bone metabolism, and cancer progression. IL-6 induces signaling through a cell surface heterodimeric receptor complex composed of a ligand binding subunit (IL-6 R alpha) and a signal transducing subunit (gp130). IL-6 binds to IL-6 R alpha, triggering IL-6 R alpha association with gp130 and gp130 dimerization. Soluble forms of IL-6 R alpha are generated by both alternative splicing and proteolytic cleavage. In a mechanism known as trans-signaling, complexes of soluble IL-6 R alpha elicit responses from

### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

# **Recombinant Human IL-6 Protein (His Tag)**

Catalog Number:PDEH100050



gp130-expressing cells that lack cell surface IL-6 R alpha. Trans-signaling enables a wider range of cell types to respond to IL-6, as the expression of gp130 is ubiquitous, while that of IL-6 R alpha is predominantly restricted to hepatocytes, monocytes, and resting lymphocytes. Soluble splice forms of gp130 block trans-signaling from IL-6/IL-6 R alpha but not from other cytokines that use gp130 as a co-receptor. IL-6, along with TNF-alpha and IL-1, drives the acute inflammatory response and the transition from acute inflammation to either acquired immunity or chronic inflammatory disease. When dysregulated, it contributes to chronic inflammation in obesity, insulin resistance, inflammatory bowel disease, arthritis, sepsis, and atherosclerosis. IL-6 can also function as an anti-inflammatory molecule, as in skeletal muscle where it is secreted in response to exercise.

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com