

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



# Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



## Recombinant Human AZGP1 protein (His tag)

Catalog Number:PDEH100173



Note: Centrifuge before opening to ensure complete recovery of vial contents.

**Description** 

Synonyms Zinc-Alpha-2-Glycoprotein;Zn-Alpha-2-GP;Zn-

Alpha-2-Glycoprotein; AZGP1; ZAG; ZNGP1

Species Human
Expression Host E.coli

Sequence Asn 148-Ser 298

Accession P25311
Calculated Molecular Weight 16.5 kDa
Observed molecular weight 22 kDa
Tag N-His

**Properties** 

**Purity** > 95 % as determined by reducing SDS-PAGE.

**Endotoxin** Please contact us for more information.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### **Background**

Zinc-α-2-Glycoprotein (AZGP1) can be found in blood plasma; seminal plasma; urine; sweat; saliva; liver; and epithelial cells of various human glands. AZGP1 has been proposed in the regulation of body weight; and the melanin production by normal and malignant melanocytes. AZGP1 stimulates lipid degradation in adipocytes and causes the extensive fat losses associated with some advanced cancers. AZGP1 has been reported to stimulate lipid breakdown and may have an important role in lipid homeostasis. Mature human AZGP1 consists of one MHC class I antigen region and a C2-type Iglike domain. AZGP1 has two alternate splice forms; one shows a 66 amino acids substitution for the C-terminal 30 amino acids; the other one shows a nine Lys substitution for amino acid 151-298.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com