



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Recombinant Human GIP protein (His tag)

Catalog Number:PDEH100174



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	gastric inhibitory polypeptide;GIP;Glucose-dependent Insulinotropic Polypeptide;Incretin
Species	Human
Expression Host	E.coli
Sequence	Glu 22-Arg 153
Accession	P09681
Calculated Molecular Weight	14.41 kDa
Observed molecular weight	18 kDa
Tag	N-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Gastric inhibitory polypeptide (GIP), also known as the glucose-dependent insulinotropic peptide is a member of the secretin family of hormones. GIP, together with glucagon-like peptide-1 (GLP-1), belongs to the group of metabolic hormones called incretins that stimulate a decrease in blood glucose levels. GIP is derived from a 153-amino acid protein encoded by the GIP gene and circulates as a biologically active 42-amino acid peptide. Engagement of Gastric inhibitory polypeptide receptors (GIPR) by GIP on pancreatic beta cells activates adenylate cyclase to regulate insulin compensation in the presence of high circulating glucose.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017