

# Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

## Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

## SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

# Recombinant Human IL-13 protein (SUMO, His tag)

Catalog Number:PDEH100188



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description				
Synonyms	ALRH;IL-13;P600;IL13			
Species	Human			
Expression Host	E.coli			
Sequence	Leu 25-Asn 146			
Accession	P35225			
Calculated Molecular Weight	33.3 kDa			
Observed molecular weight	35 kDa			
Tag	N-SUMO-His			
Properties				
Purity	> 95 % as determined by reducing SDS-PAGE.			
Endotoxin	Please contact us for more information.			
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.			
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.			
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.			
Reconstitution	Please refer to the printed manual for detailed information.			
Data				

KDa	М		R	
135	=	-		
100	-			
75				
65	•	*		
45	-			
35	-		-	
25				
15				
15				

> 95 % as determined by reducing SDS-PAGE.

#### Background

Interleukin 13 (IL-13) is a single-chain glycosylated polypeptide, which belongs to the IL-13/IL-4 family. IL-13 protein is secreted by many cell types, but especially by T helper type 2 (Th2) cells. IL-13 exerts its effects through a multi-subunit receptor comprising the alpha chain of the IL-4 receptor (IL-4R $\alpha$ ) and at least one of two known IL-13-specific binding chains (IL-13 R $\alpha$ 1 and IL-13 R $\alpha$ 2). As a cytokine, IL-13 protein is critical in regulating inflammatory, immune responses and diseases. In addition, it inhibits the production of pro-inflammatory cytokines and chemokines, and thus down-regulates macrophage activity. IL-13 protein and antibody is more importantly implicated as a central mediator of immunoregulatory processes in various cell types.

### For Research Use Only

A Reliable Research Partner in Life Science and Medicine