

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Recombinant Human ErbB4/HER4 protein (His tag)

Catalog Number:PDEH100284



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Receptor tyrosine-protein kinase erbB-4;Proto-oncogene-like protein c-

ErbB-4;Tyrosine kinase-type cell surface receptor HER4;p180erbB4

Species Human
Expression Host E.coli

Sequence Gln 26-Thr 284

AccessionQ15303Calculated Molecular Weight28.4 kDaObserved molecular weight35 kDaTagN-His

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin Please contact us for more information.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Background

ErbB4 is a type I membrane glycoprotein that is a member of the ErbB family of tyrosine kinase receptors. ErbB family members serve as receptors for the epidermal growth factor (EGF) family of growth factors. ErbB4 is expressed in normal skeletal muscle, heart, pituitary, brain and several breast carcinomas. ErbB4 ligands include the neuregulins, beta-cellulin and heparin-binding EGFlike growth factor (HBEGF). Monomeric ErbB4 binds its ligands with low affinity. Several ErbB4 isoforms exist. Two of these differ in the presence of juxtamembrane extracellular sequences which regulate the ability of TACE (TNF α converting enzyme) to proteolytically cleave ErbB4 from the cell surface. These isoforms exhibit tissuespecific expression. Another isoform lacks the phosphoinositide 3kinase activation sequence present in the ErbB4 cytoplasmic domain. ErbB4 appears to play important roles in neuronal development, development of the heart and cancer.

For Research Use Only

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com