



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Recombinant Human CD147/BSG/Basigin protein (His tag)



Catalog Number:PDEH100300

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Basigin;5F7;Collagenase Stimulatory Factor;Extracellular Matrix Metalloproteinase Inducer;EMMPRIN;Leukocyte Activation Antigen M6;OK Blood Group Antigen;Tumor Cell-Derived Collagenase Stimulatory Factor;TCSF;CD147;BSG
Species	Human
Expression Host	E.coli
Sequence	Glu 138-Ala 323
Accession	P35613
Calculated Molecular Weight	20.4 kDa
Observed molecular weight	25 kDa
Tag	N-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

CD147/EMMPRIN (Extracellular Matrix Metalloproteinase Inducer); also known as Basigin (BSG); is a transmembrane glycoprotein with different forms resulted from different modes of glycosylation and N-terminal sequence variants. It is a member of the immunoglobulin superfamily with homology to both the immunoglobulin V domain and MHC class II antigen beta-chain. This protein play important roles in variety of events including spermatogenesis; embryo implantation; neural network formation. CD147 induces the production and release of matrix metalloproteinases (MMP) in the surrounding mesenchymal cells and tumor cells; and thereby promotes invasion; metastasis; growth and survival of

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Human CD147/BSG/Basigin protein (His tag)



Catalog Number: PDEH100300

malignant cells. Furthermore; CD147 also serves as a receptor for extracellular cyclophilin and its association with integrins might be important in signal transduction. Recently; CD147 displays increased expression in many cancers; and it has been previously demonstrated to participate in cancer metastasis and progression. Thus; CD147 and its antibody are used as an effective treatment for malignant cancers.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com