

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



Recombinant Human Ataxin-3/ATXN3 protein (His tag)

Catalog Number:PDEH100314



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms Ataxin-3;ATXN3;Machado-Joseph disease protein1;Spinocerebellar ataxia type 3

proteir

SpeciesHumanExpression HostE.coli

SequenceMet 1-Lys 370AccessionP54252-2Calculated Molecular Weight40.6 kDaObserved molecular weight50 kDa

Tag N-His & C-His

Properties

Purity > 95 % as determined by reducing SDS-PAGE.

Endotoxin Please contact us for more information.

Storage Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

Shipping This product is provided as lyophilized powder which is shipped with ice packs.

Formulation Lyophilized from sterile PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

Reconstitution Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Ataxin-3 is a ubiquitously expressed nuclear protein that serves as a deubiquitinating enzyme (DUB) involved in protein homeostasis and quality control, myogenesis and degradation of misfolded chaperone substrates. As a DUB, it has two distinct features. Its N-terminal Josephin domain confers cysteine protease activity to Ataxin-3 which is important for hydrolyzing ubiquitin (Ub) linkages. The second important feature is its three Ub-interacting motifs (UIMs) through which it binds Ub conjugates and ubiquitinated proteins and bring them into proximity to trim or edit specific linkages within these Ub conjugates. It binds long poly-Ub chains and trims them. However, it has weak or no activity against chains of 4 or less Ub.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com