

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com


Recombinant Human SEC13 protein (His tag)

Catalog Number:PDEH100331



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	Protein SEC13 homolog;SEC13;GATOR complex protein SEC13;SEC13-like protein 1;SEC13-related protein
Species	Human
Expression Host	E.coli
Sequence	Met 1-Gln 322
Accession	P55735
Calculated Molecular Weight	35.3 kDa
Observed molecular weight	40 kDa
Tag	N-His & C-His
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	



> 95 % as determined by reducing SDS-PAGE.

Background

Sec13 is a ubiquitously expressed protein which participates in the formation of vesicles in the COPII complex with Sec23p.Sec24p, Sar1p and Sec31. The COPII vesicle budding complex was first described in the yeast system and was shown to provide coat proteins in the early secretory pathway. Sec13 is recruited to endoplasmic reticulum (ER) membranes where it forms a three dimensional cuboctahedron cage lattice structure by association with Sec31. This vesicular structure is necessary for ER-Golgi transport.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com