



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Recombinant Human MMP19 protein (His tag)

Catalog Number:PDEH100383



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms	Matrix metalloproteinase-19;MMP19;
Species	Human
Expression Host	E.coli
Sequence	Leu 101-Arg 450
Accession	Q99542
Calculated Molecular Weight	38.4 kDa
Observed molecular weight	41 kDa
Tag	N-His & C-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Background

MMP-19 (Matrix metalloprotease 19; also MMP-18 and MMP RASI) is a 55-59 kDa member of the peptidase M10A family of enzymes. It is widely expressed, being secreted by stratum basale keratinocytes, smooth muscle cells, epiphyseal cartilage chondrocytes and monocytes/macrophages. MMP-19 has multiple substrates, including components of the basement membrane (type IV collagen; laminin; nidogen), fibronectin, aggrecan plus COMP, and IGF-BP3, this latter cleavage resulting in the release of active IGF-I. Studies involving MMP-19 demonstrate an antiangiogenic function. This is attributable to the processing of plasminogen, generating angiostatin-like molecules, and the creation of an environment that promotes the ECM retention of soluble VEGF.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017