



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Recombinant Mouse IGFBP-2 / IGFBP2 protein (His tag)



Catalog Number:PDEM100098

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Insulin-like growth factor-binding protein 2; Igfbp2;IBP-2;IGF-binding protein 2;IGFBP-2;mIGFBP-2
Species	Mouse
Expression Host	E.coli
Sequence	Glu 35-Gln 305
Accession	P47877
Calculated Molecular Weight	29.7 kDa
Observed molecular weight	39 kDa
Tag	N-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

IGFBP-2, also known as IGFBP2, is an insulin-like growth factor-binding protein (IGFBP). IGFBPs prolong the half-life of the IGFs, control bioavailability, activity, and distribution of insulin-like growth factor (IGF) through high-affinity IGFBP/IGF complexes. Six high-affinity IGF-binding proteins (IGFBP-1 to -6) have been identified. The six IGFBPs are structurally related but encoded by distinct genes. IGFBPs have a high affinity for IGFs. Some members of the IGFBP family have been consistently shown to inhibit IGF actions by preventing them from gaining access to the IGF receptors, while others potentiate IGF actions by facilitating the ligand-receptor interaction. IGFBP-2 is overexpressed in many malignancies and is often correlated with an increasingly malignant status of the tumor, pointing to a potential

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Mouse IGFBP-2 / IGFBP2 protein (His tag)



Catalog Number: PDEM100098

involvement of IGFBP-2 in tumorigenesis. It contains 1 IGFBP N-terminal domain and 1 thyroglobulin type-1 domain. It inhibits IGF-mediated growth and developmental rates.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com