

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

Recombinant Porcine S100A12 protein (His tag)

Catalog Number: PDEP100005



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description	
Synonyms	Protein S100-A12;Calcium-binding protein in amniotic fluid 1;Calgranulin- C;Extracellular newly identified RAGE-binding protein;Migration inhibitory factor- related protein 6;S100 calcium-binding protein A12;Calcitermin;S100A12;CGRP;MRP-6;EN-RAGE
Species	Porcine
Expression Host	E.coli
Sequence	Thr 2-Glu 92
Accession	P80310
Calculated Molecular Weight	9.9 kDa
Observed molecular weight	12 kDa
Tag	N-His
Properties	
Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.
Data	



> 95 % as determined by reducing SDS-PAGE.

Background

S100 protein is a family of low molecular weight protein found in vertebrates characterized by two EF-hand calciumbinding motifs. There are at least 21 different S100 proteins, and the name is derived from the fact that the protein is 100% soluble in ammonium sulfate at neutral pH. Most S100 proteins are disulfide-linked homodimer, and is normally present in cells derived from the neural crest, chondrocytes, macrophages, dendritic cells, etc. S100 proteins have been implicated in a variety of intracellular and extracellular functions. They are involved in regulation of protein

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Catalog Number:PDEP100005



phosphorylation, transcription factors, the dynamics of cytoskeleton constituents, enzyme activities, cell growth and differentiation, and the inflammatory response. Protein S100-A12, also known as S100 calcium-binding protein A12, Calcium-binding protein in amniotic fluid 1, Calgranulin-C, and S100A12, is a member of the S-101 family. Like the majority of S100 proteins, S100A12 is a dimer, with the interface between the two subunits being composed mostly of hydrophobic residues. The fold of S100A12 is similar to the other known crystal and solution structures of S100 proteins, except for the linker region between the two EF-hand motifs. S100A12 plays an important role in the inflammatory response.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com