



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Recombinant Rat IGFBP1/IGFBP-1 protein (His tag)

Catalog Number: PDER100087



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Insulin-like growth factor-binding protein 1; Igfbp1; IBP-1; IGF-binding protein 1; IGFBP-1
Species	Rat
Expression Host	E.coli
Sequence	Ala 26-Asn 272
Accession	P21743
Calculated Molecular Weight	27.1 kDa
Observed molecular weight	35 kDa
Tag	N-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

The insulin-like growth factor binding protein (IGFBP) family consists of six structurally related proteins that bind IGF with high affinity. These proteins share conserved cysteine-rich N- and C- terminal regions that participate in IGF binding. IGFBPs regulate the bioavailability of IGFs and modulate their biological activities, both positively and negatively. Some IGFBPs also have intrinsic bioactivity that is IGF-independent. Post-translational modifications of the IGFBPs, including glycosylation, phosphorylation and proteolysis, influence IGF binding affinities and tissue localization, affecting both the IGF-dependent and independent functions.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017