

# Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



## Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

# Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in



# Recombinant Human NCR2/NKp44/CD336 Protein (His Tag)



#### Catalog Number:PDMH100003

Note: Centrifuge before opening to ensure complete recovery of vial contents.

#### **Description**

**Synonyms** CD336;dJ149M18.1;LY95;NK-p44;NKP44;RP1-149M18.2

Species Human

**Expression Host** HEK293 Cells **Sequence** Met 1-Pro 190

AccessionO95944Calculated Molecular Weight20.0 kDaObserved molecular weight37 kDaTagC-His

#### **Properties**

**Purity** > 88 % as determined by reducing SDS-PAGE.

**Endotoxin** < 1.0 EU per µg of the protein as determined by the LAL method.

**Storage** Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to

-80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots

of reconstituted samples are stable at < -20°C for 3 months.

**Shipping** This product is provided as lyophilized powder which is shipped with ice packs.

**Formulation** Lyophilized from sterile PBS, pH 7.4.


Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as

protectants before lyophilization.

Please refer to the specific buffer information in the printed manual.

**Reconstitution** Please refer to the printed manual for detailed information.

#### Data



> 88 % as determined by reducing SDS-PAGE.

#### **Background**

Natural cytotoxicity triggering receptor 2 (NCR2), also known as Natural killer cell p44-related protein (NKp44), or CD336, is a member of the natural cytotoxicity receptor (NCR) family, which composed of one Ig-like extracellular domain, a transmembrane segment, and a cytoplasmic domain. It is a novel transmembrane glycoprotein belonging to the Immunoglobulin superfamily characterized by a single extracellular V-type domain. The cytoplasmic domain of NKp44 also contains a sequence that matches the immunoreceptor tyrosine-based inhibitory motif (ITIM) consensus. This Cytotoxicity-activating receptor that may contribute to the increased efficiency of activated natural killer (NK) cells to mediate tumor cell lysis. NKp44 is selectively expressed by IL-2-activated NK cells and may contribute to the increased efficiency of activated NK cells to mediate tumor cell lysis. Tumor cell recognition of the mutated NKp44 proteins was

#### For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Fax: 1-832-243-6017

Web: www.elabscience.com Email: techsupport@elabscience.com

# Recombinant Human NCR2/NKp44/CD336 Protein (His 🍩 Tag)

Fax: 1-832-243-6017

Catalog Number:PDMH100003

significantly reduced and correlated with their lower recognition of heparin.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com