

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH


Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

Recombinant Human B7-H4/VTCN1 Protein (aa 29-258, His Tag)

Catalog Number:PDMH100005

Note: Centrifuge before opening to ensure complete recovery of vial contents.

| Description | |
|-----------------------------|---|
| Synonyms | B7S1;B7x;Vtcn1;B7h.5;B7-H4;B7H4T-cell costimulatory molecule B7x;B7S1VCTN1;B7XPRO1291;FLJ22418;Immune costimulatory protein B7-H4;Protein B7S1;T cell costimulatory molecule B7x;V-set domain containing T cell activation inhibitor 1;V-set domain-containing T-cell activation inhibitor 1 |
| Species | Human |
| Expression Host | HEK293 Cells |
| Sequence | Phe29-Ala258 |
| Accession | Q7Z7D3 |
| Calculated Molecular Weight | 26.8 kDa |
| Observed molecular weight | 43-48 kDa |
| Tag | C-His |
| Properties | |
| Purity | > 95 % as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per μ g of the protein as determined by the LAL method. |
| Storage | Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual. |
| Reconstitution | Please refer to the printed manual for detailed information. |
| Data | |



> 95 % as determined by reducing SDS-PAGE.

Background

V-set domain-containing T-cell activation inhibitor 1; also known as B7X; B7H4; B7S1; and VTCN1; is a single-pass type? membrane protein belonging to the B7 family of costimulatory proteins. These proteins are expressed on the surface of antigen-presenting cells and interact with ligands on T lymphocytes. They provide costimulatory signals that regulate T cell responses. A soluble form of B7H4 has also been detected. B7X / VTCN1 / B7H4 negatively regulates T-cell-mediated immune response by inhibiting T-cell activation; proliferation; cytokine production and development of

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

by Elabscience

Recombinant Human B7-H4/VTCN1 Protein (aa 29-258, ^{(IIII}) + A (IIII) * A (IIII) * A (IIIII) * A (IIII) *

Catalog Number:PDMH100005

cytotoxicity. When expressed on the cell surface of tumor macrophages; B7X / VTCN1 / B7H4 plays an important role; together with regulatory T-cells(Treg); in the suppression of tumor-associated antigen-specific T-cell immunity. B7X / VTCN1 / B7H4 is also involved in promoting epithelial cell transformation. This membrane protein can be up-regulated by IL6 / interleukin-6 and IL10 / interleukin-10 and inhibited by CSF2 / GM-CSF and IL4 / interleukin-4 on antigen-presenting cells.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com by Elabscience