



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Recombinant Human Frataxin/FXN protein (His tag)



Catalog Number:PDMH100050

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms	Frataxin;mitochondrial;Friedreich ataxia protein;FXN;FRDA;X25;Frataxin intermediate form
Species	Human
Expression Host	HEK293 Cells
Sequence	Leu42-Ala210
Accession	Q16595
Calculated Molecular Weight	18.5 kDa
Observed molecular weight	21 kDa
Tag	C-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Background

Frataxin intermediate form can cleave by mitochondrial processing peptidase(MPP) to the mature frataxin. Frataxin is a nuclear-encoded mitochondrial protein highly conserved in prokaryotes and eukaryotes. Its deficiency was initially described as the phenotype of Friedreich's ataxia, an autosomal recessive disease in humans. Although several functions have been described for frataxin, that is, involvement in Fe-S cluster and heme synthesis, energy conversion and oxidative phosphorylation, iron handling and response to oxidative damage, its precise function remains unclear. Although there is a general consensus on the participation of frataxin in the maintenance of cellular iron homeostasis and in iron metabolism, this protein may have other specific functions in different tissues and organisms.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017