

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

Recombinant Human TFF3 protein (His tag)

Catalog Number:PDMH100075



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Synonyms	Trefoil factor 3;Intestinal trefoil factor;hITF;Polypeptide P1.B;hP1.B;TFF3;ITF;TFI	
Species	Human	
Expression Host	HEK293 Cells	
Sequence	Met1-Phe80	
Accession	Q07654	
Calculated Molecular Weight	8.7 kDa	
Observed molecular weight	12 kDa	
Tag	C-His	
Properties		
Purity	> 95 % as determined by reducing SDS-PAGE.	
Endotoxin	Please contact us for more information.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	
Data		

KDa	М	R
80 60	11	
40	-	
30	-	8
20	-	
12	-	-

> 95 % as determined by reducing SDS-PAGE.

Background

Trefoil Factor 3 (TFF3), also known as Intestinal Trefoil Factor (ITF) and P1.B, is one of three structurally related secreted proteins that contain trefoil domains. These domains adopt a three-leaved conformation held together by conserved intrachain disulfide bonds. TFF3 is an approximately 7 kDa peptide that plays an important role in epithelial regeneration and wound healing. It can form disulfide-linked dimers or associate into disulfide-linked complexes with the intestinal mucous proteins FCGBP and MUC-2. TFF3 is expressed by epithelial goblet cells in the respiratory tract, biliary and breast ducts, small and large intestine, and cardia of the stomach. Following secretion, TFF3 can be retained in the overlying mucous layer. TFF3 is also expressed by chondrocytes during bone development. Mature human TFF3 shares 76% amino acid sequence identity with mouse and rat TFF3. TFF3 is up-regulated in response to a range of

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Catalog Number:PDMH100075



gastrointestinal epithelial disruptions. It promotes epithelial wound healing by inducing the migration of biliary, bronchial, and intestinal epithelial cells. TFF3 up-regulation is associated with and enhances tumor cell invasion and metastasis. It supports hypoxia-induced VEGF up-regulation in tumor cells and also promotes angiogenesis in non-tumor environments. Over-expression of TFF3 in type 2 diabetic mouse liver has been shown to improve glucose tolerance and insulin sensitivity.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine Toll-free: 1-888-852-8623 Tel: 1-832-243-6086 Web: www.elabscience.com Email: techsupport@elabscience.com