

Produktinformation



Forschungsprodukte & Biochemikalien
Zellkultur & Verbrauchsmaterial
Diagnostik & molekulare Diagnostik
Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien T. +43(0)1 489 3961-0 F. +43(0)1 489 3961-7 <u>mail@szabo-scandic.com</u> www.szabo-scandic.com

Recombinant Human ORM2 protein (His tag)

Catalog Number:PDMH100097



Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description		
Synonyms	AGP-B;AGP-B';AGP2;ORM2;Orosomucoid-2;OMD 2	
Species	Human	
Expression Host	HEK293 Cells	
Sequence	Met1-Ser201	
Accession	P19652	
Calculated Molecular Weight	22 kDa	
Observed molecular weight	36 kDa	
Tag	C-His	
Properties		
Purity	> 95 % as determined by reducing SDS-PAGE.	
Endotoxin	Please contact us for more information.	
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.	
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.	
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.	
Reconstitution	Please refer to the printed manual for detailed information.	
Data		

Data

kDa	MK	R
120 90 60		E.
40	-	
30		1000
20	-	•1
14	-	-
	100000000000000000000000000000000000000	

> 95 % as determined by reducing SDS-PAGE.

Background

Alpha-1-acid glycoprotein 2 is a secreted protein that belongs to the calycin superfamily; lipocalin family. It is expressed by the liver and secreted in plasma. It appears to function in modulating the activity of the immune system during the acute-phase reaction. It functions as transport protein in the blood stream. It binds various hydrophobic ligands in the interior of its beta-barrel domain. It also binds synthetic drugs and influences their distribution and availability.

For Research Use Only

A Reliable Research Partner in Life Science and MedicineToll-free: 1-888-852-8623Tel: 1-832-243-6086Web: www.elabscience.comEmail: techsupport@elabscience.com