



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Recombinant Human CD32b/FCGR2B protein (His tag)



Catalog Number:PDMH100113

**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Description

<b>Synonyms</b>	Low Affinity Immunoglobulin Gamma Fc Region Receptor II-b;IgG Fc Receptor II-b;CDw32;Fc-Gamma RII-b;Fc-Gamma-RIIb;FcRII-b;CD32;FCGR2B;FCG2;IGFR2
<b>Species</b>	Human
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Thr43-Pro217
<b>Accession</b>	P31994
<b>Calculated Molecular Weight</b>	19.1 kDa
<b>Observed molecular weight</b>	30 kDa
<b>Tag</b>	C-His

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Background

FcγRIIB is a low affinity receptor that recognizes the Fc portion of IgG. The human CD32 group consists of FcγRIIA, FcγRIIB, and FcγRIIC proteins that share 94-99% sequence identity in their extracellular domains but differ substantially in their transmembrane and cytoplasmic domains. FcγRII protein is expressed on cells of both myeloid and lymphoid lineages as well as on cells of non-hematopoietic origin. FcγRIIB has an intrinsic cytoplasmic immunoreceptor tyrosine-based inhibitory motif (ITIM) and delivers an inhibitory signal upon ligand binding. Ligation of FcγRIIB on B cells down-regulates antibody production and in some circumstances may promote apoptosis. Co-ligation of FcγRIIB on dendritic cells inhibits maturation and blocks cell activation. FcγRIIB may also be a target for monoclonal antibody therapy for malignancies. FcγRIIB plays an important negative-regulating role through modulating the signals from activation receptors.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: [www.elabscience.com](http://www.elabscience.com)

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)