



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Recombinant Human LILRB5/CD85c protein (His tag)



Catalog Number:PDMH100136

Note: Centrifuge before opening to ensure complete recovery of vial contents.

Description

Synonyms	Leukocyte immunoglobulin-like receptor subfamily B member 5;CD85 antigen-like family member C;Leukocyte immunoglobulin-like receptor 8;LIR-8;CD85c;LILRB5;LIR8
Species	Human
Expression Host	HEK293 Cells
Sequence	Met1-Gly458
Accession	O75023
Calculated Molecular Weight	50.3 kDa
Observed molecular weight	70 kDa
Tag	C-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Background

Human Leukocyte Immunoglobulin-like Receptor Subfamily B Member 5 (LILRB5/CD85c/LIR-8) belongs to a family of transmembrane glycoproteins that negatively regulate immune cell activation. Mature human LIR-8 consists of a 435 amino acid (aa) extracellular domain with four Iglike domains, a 21 aa transmembrane segment, and a 111 aa cytoplasmic domain with two immunoreceptor tyrosine-based inhibitory motifs (ITIM). Alternative splicing of human LIR-8 generates an isoform that lacks the second Ig-like domain. LIR-8 is expressed on NK cells and in the tryptic granules of mast cells. Following cell activation and degranulation, it is present on the mast cell surface. Activated mast cells may also release soluble forms of LIR-8.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017