



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Recombinant Human CD80 protein (His tag)

Catalog Number:PDMH100154



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	CD80;Activation B7-1 antigen;B7;BB1;CD28LG1;CD28LGB7-1 antigen;T-lymphocyte activation antigen CD80;B7-1;B7.1;CD28LG;LAB7
Species	Human
Expression Host	HEK293 Cells
Sequence	Met1-Asn242
Accession	P33681
Calculated Molecular Weight	26.5 kDa
Observed molecular weight	40 kDa
Tag	C-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Cluster of Differentiation 80; also called B7-1; is a member of cell surface immunoglobulin superfamily which plays key, yet distinct roles in the activation of T cells. It is the ligand for two different proteins on the T cell surface: CD28 and CTLA-4. Studies have shown that CTLA-4 binds mostly to CD80. The structure presents two extracellular domains: a membrane distal variable-like domain (IgV) and a membrane proximal Ig constant-like domain (IgC) along with an intracellular domain. Both IgV and IgC consist of anti-parallel beta sandwiches joined by a short linker region. CD80 is mostly expressed on the surface of antigen-presenting cells including activated B cells; macrophages and dendritic cells.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017