



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Recombinant Mouse B2M/Beta-2-microglobulin Protein (His Tag)



Catalog Number:PDMM100015

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	B2MG;Beta-2-Microglobulin;Beta 2 Microglobin;Beta 2 microglobulin Precursor;Beta Chain of MHC Class 1 Proteins;Beta Chain of MHC class I molecules;CDABP0092;Hdcma22p;IMD43
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met 1-Met 119
Accession	P01887
Calculated Molecular Weight	10.8 kDa
Observed molecular weight	12 kDa
Tag	C-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

B2M, also known as β 2-Microglobulin or CDABP0092, is a component of MHC class I molecules found expression in all nucleated cells (excludes red blood cells). The major function of MHC class I molecules is to display fragments of proteins from within the cell to T-cells and cells containing foreign proteins will be attacked. B2M (β 2-Microglobulin) is a low molecular weight protein. It was demonstrated that B2M (β 2-Microglobulin) was localized in the membranes of nucleated cells and was found to be associated with HL-A antigens. B2M (β 2-Microglobulin) is present in free form in various body fluids and as a subunit of histocompatibility antigens on cell surfaces lateral to the α 3 chain. Unlike α 3, β 2

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Mouse B2M/Beta-2-microglobulin Protein (His Tag)



Catalog Number:PDMM100015

has no transmembrane region. Directly above $\beta 2$ lies the $\alpha 1$ chain, which itself is lateral to the $\alpha 2$. In the absence of B2M ($\beta 2$ microglobulin), very limited amounts of MHC class I (classical and non-classical) molecules can be detected on the surface. In the absence of MHC class I, CD8 T cells, a subset of T cells involved in the development of acquired immunity cannot develop. Low levels of B2M ($\beta 2$ microglobulin) can indicate non-progression of HIV.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com