



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Recombinant Mouse Syndecan-4/Sdc4 protein (His tag)



Catalog Number:PDMM100020

Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	AA959608;AW108331;ryudocan;Synd4;syndecan-4;SDC4;SYND4;Ryudocan core protein
Species	Mouse
Expression Host	HEK293 Cells
Sequence	Met 1-Val 146
Accession	O35988
Calculated Molecular Weight	16.0 kDa
Observed molecular weight	18 kDa
Tag	C-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

SDC4 (Syndecan-4), also known as Syn4, is a transmembrane heparan sulfate proteoglycan that co-operates with integrins during cell-matrix interactions for the assembly of focal adhesions and actin stress fibers and in the phosphorylation of focal adhesion kinase (FAK) on Tyr397. Syndecan-4 plays roles in the formation of focal adhesions and stress fibers. The cytoplasmic domain of syndecan-4 interacts with a number of signalling and structural proteins, and both extracellular and cytoplasmic domains are necessary for regulated activation of associated transmembrane receptors. Syndecan-4/SDC4 is a heparan sulfate proteoglycan and works as a coreceptor for various growth factors. SDC4 deficiency limits neointimal formation after vascular injury by regulating vascular smooth muscle cells (VSMCs) proliferation and vascular progenitor

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Mouse Syndecan-4/Sdc4 protein (His tag)



Catalog Number:PDMM100020

cells (VPCs) mobilization. Therefore, SDC4 may be a novel therapeutic target for preventing arterial restenosis after angioplasty.

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com