



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

# Recombinant Rat CD163 protein (His tag)

Catalog Number: PDMR100008



**Note:** Centrifuge before opening to ensure complete recovery of vial contents.

## Description

<b>Synonyms</b>	CD163 molecule;Cd163
<b>Species</b>	Rat
<b>Expression Host</b>	HEK293 Cells
<b>Sequence</b>	Pro693-Asp939
<b>Accession</b>	A0A0G2JY2
<b>Calculated Molecular Weight</b>	27.1 kDa
<b>Observed molecular weight</b>	45 kDa
<b>Tag</b>	C-His

## Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	Please contact us for more information.
<b>Storage</b>	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

## Background

CD163, also known as M130 and p155, is a transmembrane scavenger receptor that is expressed on monocytes and macrophages and is inducible by immunosuppressant glucocorticoids and IL-10. A soluble form is shed from the cell surface by TACE or neutrophil elastase mediated cleavage in response to oxidative stress, Prostaglandin F2a stimulation, or the activation of Fc gamma receptors, TLR1, 2, 5, or 6. CD163 mediates monocyte binding to bacteria, leading to the release of inflammatory cytokines. It is essential for the circulatory clearance of hemoglobin-haptoglobin (Hb-Hp) complexes as well as free hemoglobin. It can also mediate monocyte-erythroblast adhesion and promote erythroblast expansion. CD163 binds and internalizes the cytokine TWEAK, and the ratio of soluble CD163 to TWEAK in the plasma is elevated during atherosclerosis.

## For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: [www.elabscience.com](http://www.elabscience.com)

Tel: 1-832-243-6086

Email: [techsupport@elabscience.com](mailto:techsupport@elabscience.com)

Fax: 1-832-243-6017