



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Recombinant Rat Lp-PLA2/Pla2g7 protein (His tag)

Catalog Number: PDMR100018



Note: Centrifuge before opening to ensure complete recovery of vial contents.


Description

Synonyms	Platelet-activating factor acetylhydrolase;Pla2g7;EC:3.1.1.47;rCG_43740
Species	Rat
Expression Host	HEK293 Cells
Sequence	Met1-Asn440
Accession	Q5M7T7
Calculated Molecular Weight	48.3 kDa
Observed molecular weight	60 kDa
Tag	C-His

Properties

Purity	> 95 % as determined by reducing SDS-PAGE.
Endotoxin	Please contact us for more information.
Storage	Generally, lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01 % Tween80 are added as protectants before lyophilization. Please refer to the specific buffer information in the printed manual.
Reconstitution	Please refer to the printed manual for detailed information.

Data



> 95 % as determined by reducing SDS-PAGE.

Background

Secretory phospholipase A2 is an enzyme that hydrolyses the Sn-2 ester bond of phospholipids, generating free fatty acids and lysophospholipids. Most secretory PLA2s are stored in cytoplasmic granules and are released in the extracellular environment on appropriate cell activation. Thus, they are present at higher concentration in the plasma and biologic fluids of patients with systemic inflammatory, autoimmune, or allergic disease, such as acute pancreatitis, rheumatoid arthritis, bronchial asthma, and allergic rhinitis. Also known as Lp-PLA2, PLA2G-VII is a plasma enzyme bound to lipoproteins: 80% bound to LDL, 15%-20% to HDL, and the remainder to VLDL (4-6). It is produced in major by mature macrophages and activated platelets. In contrast to other classical sPLA2s, PLA2G?VII has poor specificity toward Sn-2 long chain fatty acids, unless heavily oxidized, and undergoes the catalysis of its substrates in the aqueous phase rather

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Web: www.elabscience.com

Tel: 1-832-243-6086

Email: techsupport@elabscience.com

Fax: 1-832-243-6017

Recombinant Rat Lp-PLA2/Pla2g7 protein (His tag)

Catalog Number: PDMR100018



than at the interfacial surface of lipids. Thus, it has high specificity for water-soluble phospholipids in plasma including oxidatively modified phospholipids and platelet-activating factor (PAF). Because of the latter activity, it is also known as PAF acetylhydrolase (PAF-AH).

For Research Use Only

A Reliable Research Partner in Life Science and Medicine

Toll-free: 1-888-852-8623

Tel: 1-832-243-6086

Fax: 1-832-243-6017

Web: www.elabscience.com

Email: techsupport@elabscience.com