



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic)

## Recombinant Human LMAN2/VIP36 Protein (Human Cells, His Tag)

Catalog No. PKSH033330

### Description

<b>Synonyms</b>	Vesicular Integral-Membrane Protein VIP36; Glycoprotein GP36b; Lectin Mannose-Binding 2; Vesicular Integral-Membrane Protein 36; VIP36; LMAN2; C5orf8
<b>Species</b>	Human
<b>Expression_host</b>	Human Cells
<b>Sequence</b>	Asp45-Arg322
<b>Accession</b>	Q12907
<b>Mol_Mass</b>	32.7 kDa
<b>AP_Mol_Mass</b>	33 kDa
<b>Tag</b>	C-6His

### Properties

<b>Purity</b>	> 95 % as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg as determined by the LAL method.
<b>Storage</b>	Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of 50mM TrisHCl,10mM GSH,pH8.0.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Background

Vesicular integral-membrane protein VIP36 is also known as Glycoprotein GP36b, Lectin mannose-binding 2, Vesicular integral-membrane protein 36, LMAN2 and C5orf8. LMAN2 is widely expressed and contains one L-type lectin-like domain. LMAN2 binds high mannose type glycoproteins and may facilitate their sorting, trafficking and quality control. LMAN2 plays a role as an intracellular lectin in the early secretory pathway. LMAN2 interacts with N-acetyl-D-galactosamine and high-mannose type glycans and may also bind to O-linked glycans. LMAN2 is also involved in the transport and sorting of glycoproteins carrying high mannose-type glycans.

## SDS-PAGE

