



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Recombinant Human SIGLEC9/CD329 Protein (His Tag)(Active)

Catalog No. PKSH033787

### Description

<b>Synonyms</b>	Sialic acid-binding Ig-like lectin 9;Siglec-9;CDw329;Protein FOAP-9;SIGLEC9
<b>Species</b>	Human
<b>Expression_host</b>	Human Cells
<b>Sequence</b>	Gln18-Gly348(Lys100Glu,Ala315Glu)
<b>Accession</b>	Q9Y336
<b>Mol_Mass</b>	36.9 kDa
<b>AP_Mol_Mass</b>	55-90 kDa
<b>Tag</b>	C-His
<b>Bio_activity</b>	Measured by the ability of the immobilized protein to support the adhesion of human red blood cells. Approximately 93% will adhere after 5-10 mins incubation at 37 °C.

### Properties

<b>Purity</b>	> 95% as determined by reducing SDS-PAGE.
<b>Endotoxin</b>	< 1.0 EU per µg as determined by the LAL method.
<b>Storage</b>	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks.Reconstituted protein solution can be stored at 4-7°C for 2-7 days.Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS?2mMEDTA?ph7.4.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Background

Sialic acid-binding Ig-like lectin 9(Siglec 9) is expressed by peripheral blood leukocytes (neutrophils and monocytes but not eosinophils), and found in liver, fetal liver, bone marrow, placenta, spleen and in lower levels in skeletal muscle, fetal brain and so on. It is a putative adhesion molecule that mediates sialic-acid dependent binding to cells. It also binds to alpha-2,3- or alpha-2,6-linked sialic acid. The sialic acid recognition site may be masked by cis interactions with sialic acids on the same cell surface.

## SDS-PAGE

