



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Recombinant Human Glypican-3/GPC3/OCI5 (C-6His-Avi) Biotinylated

Catalog No. PKSH033852

Description

| | |
|------------------------|---|
| Synonyms | Sialic acid-binding Ig-like lectin 9;Siglec-9;CDw329;Protein FOAP-9;SIGLEC9 |
| Species | Human |
| Expression_host | Human Cells |
| Sequence | Gln25-His559 |
| Accession | P51654 |
| Mol_Mass | 63.4 kDa |
| AP_Mol_Mass | 70-130&40 kDa |
| Tag | C-6His-Avi |
| Bio_activity | Immobilized Human FGFb (Cat#PKSH032439) at 10µg/ml (100 µl/well) can bind Human Glypican-3-His(Cat#PKSH033852). The ED50 of Human NRG1Beta-His (Cat#PKSH033852) is 4.5 ug/ml. |

Properties

| | |
|-----------------------|--|
| Purity | >95% as determined by reducing SDS-PAGE. |
| Endotoxin | < 1.0 EU per µg as determined by the LAL method. |
| Storage | Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months. |
| Shipping | This product is provided as lyophilized powder which is shipped with ice packs. |
| Formulation | Lyophilized from a 0.2 µm filtered solution of PBS, pH 7.4. |
| Reconstitution | Please refer to the printed manual for detailed information. |

Background

Glypican-3/GPC3 is a member of the glypican family. It belongs to the glypican family and is highly expressed in lung, liver, and kidney. It is a heparan sulfate proteoglycan, which is overexpressed in various neoplasms such as hepatocellular carcinoma, malignant melanoma, and testicular yolk sac tumor, and plays an important role in cell growth and differentiation. GPC3 function is tissue dependent. In some tissues, GPC3 acts as a tumor suppressor gene, whereas in others, it acts as an oncofetal protein. GPC3 is a reliable marker for hepatocellular carcinoma. The sensitivity and specificity exceeds both alpha-fetoprotein and hepatocyte-paraffin1. GPC3 immunohistochemistry can aid in the differentiation of testicular germ cell tumors, being expressed in all yolk sac tumors but not in seminomas. GPC3 expression has also been identified in some squamous cell carcinomas of the lung and clear cell carcinomas of the ovary. Glypican-3 is currently regarded as a tumor marker and potential target for immunotherapy.

SDS-PAGE

