



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

## Recombinant Mouse MBL2/MBP-C Protein

Catalog No. PKSM041108

### Description

<b>Synonyms</b>	Mannose binding lectin (C); isoform CRA_b; Mannose-binding protein C; Mbl2; MBL-2; Mannose Binding Lectin 2
<b>Species</b>	Mouse
<b>Expression_host</b>	Human Cells
<b>Sequence</b>	Glu19-Asp244
<b>Accession</b>	Q3UEK1
<b>Mol_Mass</b>	24.0 kDa
<b>AP_Mol_Mass</b>	28 kDa

### Properties

<b>Purity</b>	> 95 % as determined by SDS-PAGE
<b>Endotoxin</b>	< 1.0 EU per µg as determined by the LAL method.
<b>Storage</b>	Lyophilized proteins are stable for up to 12 months when stored at -20 to -80°C. Reconstituted protein solution can be stored at 4-8°C for 2-7 days. Aliquots of reconstituted samples are stable at < -20°C for 3 months.
<b>Shipping</b>	This product is provided as lyophilized powder which is shipped with ice packs.
<b>Formulation</b>	Lyophilized from a 0.2 µm filtered solution of PBS, pH7.4.
<b>Reconstitution</b>	Please refer to the printed manual for detailed information.

### Background

Mannose-binding Lectin (MBL) is an acute phase protein bearing to the family of collectins produced by the liver as a monomer that forms a triple helix. Once released in serum, it further polymerizes forming dimers to octamers. The degree of serum polymerization is critical for the biological activity of MBL. MBL has higher affinity to microbial polysaccharides or their glycoconjugates. MBL was shown earlier to bind cell surfaces of bacteria, fungi, protozoa and viruses and acts as an acute-phase plasma protein (APP) during infection and inflammation. MBL activates the lectin-complement pathway, promotes opsonophagocytosis and modulates inflammation.

## SDS-PAGE

