



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

Recombinant Cynomolgus SIGLEC5/CD170 Protein (His Tag)

Catalog No. PKSQ050010

Description

Synonyms	Sialic acid-binding Ig-like lectin 5; Siglec-5; CD170;CD33L2;OB-BP2;OBBP;OBBP2;SIGLEC-5;SIGLEC5
Species	Cynomolgus
Expression_host	Human Cells
Sequence	Glu17-Gly435
Accession	A0A0B4J1D1
Mol_Mass	46.8 kDa
AP_Mol_Mass	90 kDa
Tag	C-His

Properties

Purity	> 90% as determined by reducing SDS-PAGE.
Endotoxin	< 1.0 EU per µg as determined by the LAL method.
Storage	Lyophilized protein should be stored at < -20°C, though stable at room temperature for 3 weeks.Reconstituted protein solution can be stored at 4-7°C for 2-7 days.Aliquots of reconstituted samples are stable at < -20°C for 3 months.
Shipping	This product is provided as lyophilized powder which is shipped with ice packs.
Formulation	Lyophilized from a 0.2 µm filtered solution of PBS, pH8.0.
Reconstitution	Please refer to the printed manual for detailed information.

Background

Sialic acid-binding Ig-like lectin 5 is a protein that in Cynomolgus is encoded by the SIGLEC5 gene, Cynomolgus SIGLEC5 cDNA encodes 551 amino acids (aa) that include a 16 aa signal sequence, a 439aa extracellular domain (ECD) with three Ig-like domains, a transmembrane region and a cytoplasmic tail. No Siglec has been shown to recognize any cell surface ligand other than sialic acids, suggesting that interactions with glycans containing this carbohydrate are important in mediating the biological functions of Siglecs. Siglec5 to 11 share a high degree of sequence similarity with CD33/Siglec3 both in their extracellular and intracellular regions. Putative adhesion molecule that mediates sialic-acid dependent binding to cells. Binds equally to alpha-2,3-linked and alpha-2,6-linked sialic acid. The sialic acid recognition site may be masked by cis interactions with sialic acids on the same cell surface.

SDS-PAGE

