



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

DMEM (High glucose) (without L-glutamine)

Cat. No: PM150213
Size: 500mL

General Information

Product Form	Liquid
Concentration	1×
pH	7.2-7.4
D-glucose	4500mg/L
HEPES	Negative
L-glutamine	Negative
NaHCO ₃	3700mg/L
Phenol red	15mg/L
Sodium pyruvate	1 mM
Storage	2-8°C, Shading light
Shipping	RT
Expiration date	12 months

Background

DMEM (Dulbecco's Modified Eagle Medium) was developed on the basis of MEM medium. Compared with MEM medium, the content of amino acid increased by 2 times, the content of vitamin increased by 4 times, and the content of non-essential amino acid, trace iron ion and sodium pyruvate were increased by 4 times. The glucose content of DMEM medium was originally designed as 1000 mg/L (low Glucose type), and then developed into 4500mg/L (high Glucose type), which has been widely used in cell culture. DMEM (High glucose) was widely used in fast growth, low adhesion cells, hybridoma myeloma cells, clone cells, DNA transfected transformation cells, various primary virus host cells, single cell culture and vaccine production. DMEM (High glucose) contains many kinds of amino acids, vitamins, inorganic salts and other ingredients for cell culture, but does not contain protein, lipids or any growth factors, so the product should be used with serum or serum-free additives.

Notes

1. This product is for research use only.
2. This product is sterilized by 0.22 μ m filtration.
3. It is necessary to pay attention to the aseptic operation and avoid the pollution during the culture.