

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten! See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere Liefer- und Versandbedingungen

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

linkedin.com/company/szaboscandic in





DMEM (glucose free) (without phenol red, L-glutamine)

Cat. No: PM150279 Size: 500mL

General Information

Product Form Liquid Concentration 1xрH 7.2 - 7.4D-glucose Negative **HEPES** Negative L-glutamine Negative NaHCO₃ 3700mg/L Phenol red Negative Sodium pyruvate 1 mM

Storage 2-8°C, Shading light

Shipping RT

Expiration date 12 months

Background

DMEM (Dulbecco's Modified Eagle Medium) was developed on the basis of MEM medium. Compared with MEM medium, the content of amino acid increased by 2 times, the content of vitamin increased by 4 times, and the content of non-essential amino acid, trace iron ion and sodium pyruvate were increased by 4 times. The glucose content of DMEM medium was originally designed as 1000 mg/L (low Glucose type), and then developed into 4500mg/L (high Glucoser type), which has been widely used in cell culture. DMEM medium without glucose can be added to the concentration of glucose at will according to the research needs, which is convenient and quick. DMEM (glucose free) contains many kinds of amino acids, vitamins, inorganic salts and other ingredients for cell culture, but does not contain protein, lipids or any growth factors, so the product should be used with serum or serum-free additives.

Notes

- 1. This product is for research use only.
- 2. This product is sterilized by 0.22 µ m filtration.
- 3. It is necessary to pay attention to the aseptic operation and avoid the pollution during the culture.