



# SZABO SCANDIC

Part of Europa Biosite

## Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!  
See the following pages for more information!



### Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

### Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

### SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

[mail@szabo-scandic.com](mailto:mail@szabo-scandic.com)

[www.szabo-scandic.com](http://www.szabo-scandic.com)

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



## UK Office

### Everest Biotech Ltd

Cherwell Innovation Centre  
77 Heyford Park  
Upper Heyford  
Oxfordshire  
OX25 5HD  
UK

Enquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[sales@everestbiotech.com](mailto:sales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

## US Office

### Everest Biotech c/o Abcore

405 Maple Street, Suite A106  
Ramona,  
CA 92065  
USA

Inquiries:

[info@everestbiotech.com](mailto:info@everestbiotech.com)

Sales:

[usasales@everestbiotech.com](mailto:usasales@everestbiotech.com)

Tech support:

[support@everestbiotech.com](mailto:support@everestbiotech.com)

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

[www.everestbiotech.com](http://www.everestbiotech.com)

**Research Use Only. Not for  
diagnostic or therapeutic use.**

## EB05507 - Goat Anti-AIRE Antibody

Size: 100µg specific antibody in 200µl



### Target Protein

**Principal Names:** AIRE, autoimmune regulator (autoimmune polyendocrinopathy candidiasis ectodermal dystrophy), APS1, APSI, PGA1, APECED, AIRE protein, autoimmune regulator (APECED protein), autoimmune regulator (autoimmune polyendocrinopathy candidiasis ectodermal dystrophy), autoimmune regulator, AIRE1, OTTHUMP00000109529, autoimmune polyendocrinopathy candidiasis ectodermal dystrophy

**Official Symbol:** AIRE

**Accession Number(s):** NP\_000374.1

**Human GeneID(s):** [326](#)

### Immunogen

Peptide with sequence C-QSMARPAAPFPS, from the C Terminus of the protein sequence according to NP\_000374.1.

Please note the [peptide](#) is available for sale.

### Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

### Applications Tested

**Peptide ELISA:** antibody detection limit dilution 1:128000.

**Western blot:** Preliminary testing showed an approx 60kDa band in nuclear lysates of cell line Jurkat and in Human Spleen lysates (calculated MW of 57.7kDa according to NP\_000374.1). Recommended concentration: 1-2µg/ml. Primary incubation 1 hour at room temperature. **Negative Control:** Human Smooth Muscle lysate. This product has been successfully used in Western blot by an anonymous customer on HEK293 cell lysate.

**IHC:** In paraffin embedded Human Thymic medulla shows staining of select nuclei. Recommended concentration: 2.5µg/ml.


**Immunofluorescence:** Strong expression of the protein seen in the nucleoplasm and nuclear membranes of U2OS and HeLa cells. Recommended concentration: 10µg/ml.

**Flow Cytometry:** Flow cytometric analysis of Jurkat cells. Recommended concentration: 10ug/ml.


### Species Reactivity

**Tested:** Human


**Expected from sequence similarity:** Human




EB05507 (1 $\mu$ g/ml) staining of nuclear Jurkat cell (A), (2 $\mu$ g/ml) Human Spleen (B) and negative control Human Smooth Muscle (C) lysate (35 $\mu$ g protein in RIPA buffer). Detected by chemiluminescence.




EB05507 (2 $\mu$ g/ml) staining of paraffin embedded Human Thymus. Steamed antigen retrieval with Tris/EDTA buffer pH 9, HRP-staining.



EB05507 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nucleoplasm and nuclear membrane staining. Actin filaments were stained with phalloidin (red) and the nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB05507 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing nucleoplasm and nuclear membrane staining. Actin filaments were stained with phalloidin (red) and the nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB05507 Flow cytometric analysis of paraformaldehyde fixed Jurkat cells (blue line), permeabilized with 0.5% Triton. Primary incubation overnight (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.