



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06233 - Goat Anti-KPNA2 / IPOA1 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: KPNA2, IPOA1, RCH1, SRP1alpha, karyopherin alpha 2 (RAG cohort 1, importin alpha 1), RAG cohort 1, importin alpha 1, importin alpha 2, importin-alpha-P1, karyopherin subunit alpha-2, pendulin, importin subunit alpha-2, QIP2, RAG cohort protein 1, SRP1-alpha

Official Symbol: KPNA2

Accession Number(s): NP_002257.1

Human GeneID(s): [3838](#)

Non-Human GeneID(s): 16647 (mouse), 85245 (rat)

Immunogen

Peptide with sequence C-QVQDGAPGTFNF, from the C Terminus of the protein sequence according to NP_002257.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:32000.

Western blot: Approx 55kDa band observed in lysates of cell lines CaCo-2, Jurkat, and KNRK, and approx. 58kDa in lysates of cell lines MCF7 and A549 (calculated MW of 57.9kDa according to Human NP_002257.1 and Rat NP_001257731.1). Recommended concentration 0.03-0.1µg/ml. Primary incubation 1 hour at room temperature.

IHC: Paraffin embedded Human Testis. Recommended concentration: 5µg/ml.

Immunofluorescence: Strong expression of the protein seen in U2OS and A549 cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of A549 cells. Recommended concentration: 10µg/ml.

Species Reactivity

Tested: Human, Rat

Expected from sequence similarity: Human, Mouse, Rat, Dog

Specific References

This antibody (previous batch) has been successfully used in Western blot on Mouse:

Ma KL, Song LK, Yuan YH, Zhang Y, Han N, Gao K, Chen NH

The nuclear accumulation of alpha-synuclein is mediated by importin alpha and promotes neurotoxicity. *Neuropharmacology*. 2014 Jul;82:132-42

PMID: 23973294

This antibody has been successfully used in the following paper:


Fujimoto H, Ikuta T, Koike A and Koike M

Acetylation of nuclear localization signal controls importin-mediated nuclear transport of


Ku70

<https://doi.org/10.1101/403485> (August 29, 2018)


PMID: 0




EB06233 (0.1µg/ml) staining of Jurkat (A) and CaCo-2 (B) and (0.03ug/ml) A549 (C), MCF7 (D) and KNRK (E) cell lysate. (35µg protein in RIPA buffer). Detected by chemiluminescence.




EB06233 (5µg/ml) staining of paraffin embedded Human Testis. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.



EB06233 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic and ER/Golgi staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB06233 Immunofluorescence analysis of paraformaldehyde fixed A549 cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing cytoplasmic staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB06233 Flow cytometric analysis of paraformaldehyde fixed A549 cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.