



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 



UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06374 - Goat Anti-ECSM2 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: endothelial cell-specific marker 2, endothelial cell-specific molecule 2, hypothetical protein LOC641700

Official Symbol: ECSM2

Accession Number(s): NP_001071161.1

Immunogen

Peptide with sequence C-NNGKQSLSAEKVL, from the C Terminus of the protein sequence according to NP_001071161.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:128000.

Western blot: Approx 50kDa band observed in Human Lung and Human Umbilical Cord lysates (calculated MW of 21.3kDa according to NP_001071161.1). The observed molecular weight corresponds to earlier findings in literature with different antibodies (Ref PMID: 18556573). Recommended concentration: 1-3µg/ml.

IHC: Paraffin embedded Human Vessel. Recommended concentration: 5µg/ml. This product has been successfully used in IHC on Human: Matsuo K et al, J Biol Chem. 2015 Feb 6;290(6):3784. PMID: 25533470.

Immunofluorescence: This product has been successfully used in IF on Human: Matsuo K et al, J Biol Chem. 2015 Feb 6;290(6):3784. PMID: 25533470.

Species Reactivity

Tested: Human

Expected from sequence similarity: Human


Specific Reference

This antibody has been successfully used in IHC and IF on Human:


Matsuo K, Akakabe Y, Kitamura Y, Shimoda Y, Ono K, Ueyama T, Matoba S, Yamada H, Hatakeyama K, Asada Y, Emoto N, Ikeda K.

Loss of apoptosis regulator through modulating IAP expression (ARIA) protects blood vessels from
J Biol Chem. 2015 Feb 6;290(6):3784

PMID: 25533470



EB06374 (1µg/ml) staining of Human lung (A) and Human Umbilical Cord (B) lysates (35µg protein in RIPA buffer). Primary incubation was 1 hour.



EB06374 (5µg/ml) staining of paraffin embedded Human Vessel. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.