



SZABO SCANDIC

Part of Europa Biosite

Produktinformation



Forschungsprodukte & Biochemikalien



Zellkultur & Verbrauchsmaterial



Diagnostik & molekulare Diagnostik



Laborgeräte & Service

Weitere Information auf den folgenden Seiten!
See the following pages for more information!



Lieferung & Zahlungsart

siehe unsere [Liefer- und Versandbedingungen](#)

Zuschläge

- Mindermengenzuschlag
- Trockeneiszuschlag
- Gefahrgutzuschlag
- Expressversand

SZABO-SCANDIC HandelsgmbH

Quellenstraße 110, A-1100 Wien

T. +43(0)1 489 3961-0

F. +43(0)1 489 3961-7

mail@szabo-scandic.com

www.szabo-scandic.com

[linkedin.com/company/szaboscandic](https://www.linkedin.com/company/szaboscandic) 

UK Office

Everest Biotech Ltd

Cherwell Innovation Centre
77 Heyford Park
Upper Heyford
Oxfordshire
OX25 5HD
UK

Enquiries:

info@everestbiotech.com

Sales:

sales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: +44 (0)1869 238326

Fax: +44 (0)1869 238327

US Office

Everest Biotech c/o Abcore

405 Maple Street, Suite A106
Ramona,
CA 92065
USA

Inquiries:

info@everestbiotech.com

Sales:

usasales@everestbiotech.com

Tech support:

support@everestbiotech.com

Tel: 888-320-4628 (toll-free)

Fax: 888-841-9041

www.everestbiotech.com

**Research Use Only. Not for
diagnostic or therapeutic use.**

EB06471 - Goat Anti-EndoPDI / TXNDC5 Antibody

Size: 100µg specific antibody in 200µl



Target Protein

Principal Names: EndoPDI, thioredoxin domain containing 5, ERP46, UNQ364, MGC3178, FLJ21353, FLJ90810, thioredoxin related protein, endothelial protein disulphide isomerase, TXNDC5, thioredoxin domain containing 5 (endoplasmic reticulum), Hcc-2, PDIA15, endothelial protein disulphide isomerase, protein disulfide isomerase family A, member 15, thioredoxin domain containing 5

Official Symbol: TXNDC5

Accession Number(s): NP_110437.2; NP_001139021.1

Human GeneID(s): [81567](#)

Important Comments: This antibody is expected to be able to recognise both reported human isoforms, as represented by NP_110437.2; NP_001139021.1.

Immunogen

Peptide with sequence C-SLHRFVLSQAKDEL, from the C Terminus of the protein sequence according to NP_110437.2; NP_001139021.1.

Please note the [peptide](#) is available for sale.

Purification and Storage

Purified from goat serum by ammonium sulphate precipitation followed by antigen affinity chromatography using the immunizing peptide.

Supplied at 0.5 mg/ml in Tris saline, 0.02% sodium azide, pH7.3 with 0.5% bovine serum albumin.

Aliquot and store at -20°C. Minimize freezing and thawing.

Applications Tested

Peptide ELISA: antibody detection limit dilution 1:64000.

Western blot: Approx 50kDa band observed in lysates of cell lines HEK293, A549, HeLa and HepG2 (calculated MW of 47.6kDa according to NP_110437.2). This molecular weight is routinely observed by other sources. Recommended concentration: 0.01-0.1µg/ml. Primary incubation 1 hour at room temperature.

IHC: Paraffin embedded Human Small Intestine. Recommended concentration: 3.75µg/ml.


Immunofluorescence: Strong expression of the protein seen in the endoplasmic reticulum of U2OS and HeLa cells. Recommended concentration: 10µg/ml.

Flow Cytometry: Flow cytometric analysis of HeLa cells. Recommended concentration: 10ug/ml.


Species Reactivity

Tested: Human


Expected from sequence similarity: Human




EB06471 staining (0.1 μ g/ml) of HEK293 (A), A549 (B), HeLa (C) and (0.01 μ g/ml) of HepG2 (D) cell lysate (35 μ g protein in RIPA buffer). Detected by chemiluminescence.




EB06471 Immunofluorescence analysis of paraformaldehyde fixed HeLa cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10 μ g/ml) followed by Alexa Fluor 488 secondary antibody (2 μ g/ml), showing endoplasmic reticulum staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10 μ g/ml) followed by Alexa Fluor 488 secondary antibody (2 μ g/ml).



EB06471 Immunofluorescence analysis of paraformaldehyde fixed U2OS cells, permeabilized with 0.15% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml), showing endoplasmic reticulum staining. The nuclear stain is DAPI (blue). Negative control: Unimmunized goat IgG (10ug/ml) followed by Alexa Fluor 488 secondary antibody (2ug/ml).



EB06471 Flow cytometric analysis of paraformaldehyde fixed HeLa cells (blue line), permeabilized with 0.5% Triton. Primary incubation 1hr (10ug/ml) followed by Alexa Fluor 488 secondary antibody (1ug/ml). IgG control: Unimmunized goat IgG (black line) followed by Alexa Fluor 488 secondary antibody.



EB06471 (3.75µg/ml) staining of paraffin embedded Human Small Intestine. Steamed antigen retrieval with citrate buffer pH 6, AP-staining.